

Gibbsianness and non-Gibbsianness in divide and colour models

András Bálint *

Department of Mathematics, VU University Amsterdam

April 7, 2009

Abstract

For parameters $p \in [0, 1]$ and $q > 0$ such that the random-cluster measure $\Phi_{p,q}^{\mathbb{Z}^d}$ for $\mathbb{Z}^d$ with parameters p and q is unique, the q -divide and colour (DaC(q)) model on $\mathbb{Z}^d$ is defined as follows. First we draw a bond configuration with distribution $\Phi_{p,q}^{\mathbb{Z}^d}$. Then to each FK cluster (i.e., to every vertex in the FK cluster), independently for different FK clusters, we assign a spin from the set $\{1, 2, \dots, s\}$ in such a way that spin i has probability a_i .

In this paper we prove that the resulting measure on spin configurations is a Gibbs measure for small values of p , and is not a Gibbs measure for large p , except in the special case of $q \in \{2, 3, \dots\}$, $a_1 = a_2 = \dots = a_s = 1/q$, when the DaC(q) model coincides with the q -state Potts model.

Keywords: Gibbs measures, non-Gibbsianness, random-cluster measures, quasilocality, DaC models

AMS 2000 Subject Classification: 60K35, 82B20, 82B43

1 Introduction

The random-cluster representations of various models have played an important role in the study of physical systems and phase transitions. They provide a different viewpoint of the physical models, and many problems in the Ising and Potts models can indeed be solved by using their random-cluster representations, see e.g., [10, 16, 11].

*E-mail: abalint@few.vu.nl

For $\beta \geq 0$ and an integer $q \geq 2$, a spin configuration in the q -state Potts model at inverse temperature β can be obtained as follows. Draw a bond configuration according to a random-cluster measure with parameters $p = 1 - e^{-2\beta}$ and q (for definitions, see Section 2), then assign to each vertex a spin from the set $\{1, 2, \dots, q\}$ in such a way that all spins have equal probability and that vertices that are connected in the bond configuration get the same spin. If the spin is chosen from a set $\{1, 2, \dots, s\}$ with an integer $1 < s < q$, and the probability of spin i is k_i/q with integers $k_1, k_2, \dots, k_s$ such that $\sum_{i=1}^s k_i = q$, then we get the so-called fuzzy Potts model [23, 19]. Recent papers (see [17, 18, 9, 22, 1, 2, 12]) have shown that generalisations of the above constructions with different values of q and s , and more general rules of spin assignment are also of interest. From a mathematical viewpoint, such models are natural examples of a dependent site percolation model with a simple definition but nontrivial behaviour. The study of such models may also lead to a better understanding of models of primary physical importance, as it was the case in [2] where an informative, new perspective of the high temperature Ising model on the triangular lattice was given.

The model treated here is defined as follows. Let $G = (\mathcal{V}, \mathcal{E})$ be a (finite or infinite) locally finite graph. Fix parameters $p \in [0, 1]$, $q > 0$ in such a way that there exists exactly one random-cluster measure for G with parameters p and q . We denote this measure by $\Phi_{p,q}^G$. Fix also an integer $s \geq 2$, and $a_1, a_2, \dots, a_s \in (0, 1)$ such that $\sum_{i=1}^s a_i = 1$, and define the **single-spin space** $S = \{1, 2, \dots, s\}$, and the **state space** $\Omega^G = \Omega_C^G \times \Omega_D^G$ with $\Omega_C^G = S^\mathcal{V}$ and $\Omega_D^G = \{0, 1\}^\mathcal{E}$. Let Y be a random bond configuration taking values in Ω_D^G with distribution $\Phi_{p,q}^G$. Given $Y = \eta$ for some $\eta \in \Omega_D^G$, we construct a random Ω_C^G -valued spin configuration X by assigning spin $i \in S = \{1, 2, \dots, s\}$ with probability a_i to each connected component in η (i.e., the same spin i to each vertex in the component), independently for different components. We write $\mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^G$ for the joint distribution of (X, Y) on Ω , and $\mu_{p,q,(a_1,a_2,\dots,a_s)}^G$ for the marginal of $\mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^G$ on Ω_C^G . This definition is a slight generalisation of the fractional fuzzy Potts model defined in [17], p.1156 (see also [1], Section 1.2). However, we shall call this model the **q -divide and colour (DaC(q)) model** to emphasise that we look at it rather as a generalisation of the model introduced in [18] by Häggström (which is the DaC(1) model in the present terminology) than of the fuzzy Potts model of [23, 19].

Let us now consider the (hypercubic) lattice with vertex set $\mathbb{Z}^d$ and edge set $\mathcal{E}^d$ with edges between vertices at Euclidean distance 1. With an abuse of notation, we shall denote this graph by $\mathbb{Z}^d$, and the sets $\Omega_D^{\mathbb{Z}^d}, \Omega_C^{\mathbb{Z}^d}$ and $\Omega^{\mathbb{Z}^d}$ by Ω_D, Ω_C and Ω , respectively. The present work is focused on the Gibbs properties and k -Markovianity of the measure $\mu_{p,q,(a_1,a_2,\dots,a_s)}^{\mathbb{Z}^d}$ in $d \geq 2$

dimensions. Since the cases $p = 0$ or 1 are trivial, we henceforth assume $p \in (0, 1)$. We give results for $q \geq 1$ only, since much more is known about random-cluster measures with $q \geq 1$ than with $q < 1$.

We shall prove that, except in the special case of $q = s$ and $a_1 = a_2 = \dots = a_s$ (when the $\text{DaC}(q)$ model coincides with the q -state Potts model on $\mathbb{Z}^d$ at inverse temperature $\beta = -\frac{1}{2} \log(1 - p)$), the $\text{DaC}(q)$ model is not k -Markovian for any k . For large values of p , $\mu_{p,q,(a_1,a_2,\dots,a_s)}^{\mathbb{Z}^d}$ is not even quasilocal and is therefore not a Gibbs measure, again with the exception of the Potts case. This shows that the Gibbsianness of the Potts model at low temperatures is very sensitive with respect to perturbations in the assignment of the spin probabilities, even the smallest change makes the model non-quasilocal. Some intuition behind this phenomenon will be given after Lemma 3.3. By demonstrating the special role of the q -state Potts Gibbs measure among $\text{DaC}(q)$ models, our result supports the view expressed in [5, 6] that, especially at low temperatures, Gibbsianness of measures is rather a lucky exception than the rule. However, if p is small enough, then Gibbsianness does hold. The proof of this fact uses the idea that at small values of p (which corresponds to high temperatures in the (fuzzy) Potts model), the $\text{DaC}(q)$ model is close in spirit to independent site percolation on $\mathbb{Z}^d$.

These results are in line with those in [19] (see also [23]) concerning the fuzzy Potts model, and in some cases, essentially the same proofs work in the current, more general situation. Therefore, at places only a sketch of the proof is given, and the reader is referred to [19] for the details. Note, however, that such similarities are not at all immediate from the definition of the models. More importantly, in the $\text{DaC}(q)$ model, a distinction must be made between the case when $a_i \geq 1/q$ for all i , and when there exists j with $a_j < 1/q$. In the former (of which the fuzzy Potts model is a special case), a rather complete picture can be given, whereas in the latter, there is an interval in p where we do not know whether $\mu_{p,q,(a_1,a_2,\dots,a_s)}^{\mathbb{Z}^d}$ is a Gibbs measure.

Finally, we give a sufficient (but not necessary) condition for almost sure quasilocality of $\mu_{p,q,(a_1,a_2,\dots,a_s)}^{\mathbb{Z}^d}$, and as an application, we obtain this weak form of Gibbsianness in the two-dimensional case for a large range of parameters.

2 Definitions and main results

2.1 Random-cluster measures

In this section, we recall the definition of the Fortuin-Kasteleyn (FK) random-cluster measures, and those properties of these measures that are important

for the rest of the paper. For the proofs and much more on random-cluster measures, see, e.g., [15].

Definition 2.1. For a finite graph $G = (\mathcal{V}, \mathcal{E})$ and parameters $p \in [0, 1]$ and $q > 0$, the **random-cluster measure** $\Phi_{p,q}^G$ is the measure on Ω_D^G which assigns to a bond configuration $\eta \in \Omega_D^G$ probability

$$\Phi_{p,q}^G(\eta) = \frac{q^{k(\eta)}}{Z_{p,q}^G} \prod_{e \in \mathcal{E}} p^{\eta(e)} (1-p)^{1-\eta(e)}, \quad (1)$$

where $k(\eta)$ is the number of connected components in the graph with vertex set $\mathcal{V}$ and edge set $\{e \in \mathcal{E} : \eta(e) = 1\}$ (we call such components **FK clusters** throughout, edges with state 1 **open**, and edges with state 0 **closed**), and $Z_{p,q}^G$ is the appropriate normalising factor.

This definition is not suitable for infinite graphs. In that case, we shall require that certain conditional probabilities are the same as in the finite case. A graph is called **locally finite** if every vertex has a bounded degree. We shall denote bond configurations throughout by η and ζ . For the restriction of a bond configuration η to an edge set H , we write η_H . For vertices v and w , we denote the edge between v and w by $\langle v, w \rangle$. The following definition is taken from [19], and its equivalence with a more common definition (where arbitrary finite edge sets and not only single edges are considered) is stated e.g. in Lemma 6.18 of [11].

Definition 2.2. For an infinite, locally finite graph $G = (\mathcal{V}, \mathcal{E})$ and parameters $p \in [0, 1]$, $q > 0$, a measure ϕ on Ω_D^G is called a **random-cluster measure for G with parameters p and q** if for any edge $e = \langle x, y \rangle \in \mathcal{E}$, and edge configuration $\zeta \in \{0, 1\}^{\mathcal{E} \setminus \{e\}}$ outside e , we have that

$$\phi(\{\eta \in \Omega_D^G : \eta(e) = 1\} \mid \{\eta \in \Omega_D^G : \eta_{\mathcal{E} \setminus \{e\}} = \zeta\}) = \begin{cases} p & \text{if } x \overset{\zeta}{\leftrightarrow} y, \\ \frac{p}{p+(1-p)q} & \text{otherwise,} \end{cases}$$

where $x \overset{\zeta}{\leftrightarrow} y$ denotes that there exists a path of edges between x and y in which every edge has ζ -value 1.

It is not difficult to prove that one gets the same conditional probabilities for random-cluster measures on finite graphs, hence Definition 2.2 is a reasonable extension of Definition 2.1 to infinite graphs.

It is not clear from the definition that such measures exist. However, for $\mathbb{Z}^d$ and for $q \geq 1$, two random-cluster measures can be constructed as follows. For a vertex set $H \subset \mathbb{Z}^d$, let ∂H denote the **vertex boundary** of

the set, that is, $\partial H = \{v \in \mathbb{Z}^d \setminus H : \exists w \in H \text{ such that } \langle v, w \rangle \in \mathcal{E}^d\}$. Define, for $n = 1, 2, \dots$, the set $\Lambda_n = \{-n, \dots, n\}^d$, the graph $G_n = (\mathcal{V}_n, \mathcal{E}_n)$ with vertex set $\mathcal{V}_n = \Lambda_n \cup \partial\Lambda_n$ and edge set $\mathcal{E}_n = \{e \in \mathcal{E}^d : \text{both endvertices of } e \text{ are in } \mathcal{V}_n\}$. For $n = 1, 2, \dots$, let W_n be the event that all edges with both endvertices in $\partial\Lambda_n$ are open, and let $\Phi_{p,q}^{G_n,1}$ be the measure $\Phi_{p,q}^{G_n}$ conditioned on W_n . Then both $\Phi_{p,q}^{G_n}$ and $\Phi_{p,q}^{G_n,1}$ converge weakly as $n \rightarrow \infty$; we denote the limiting measures by $\Phi_{p,q}^{\mathbb{Z}^d,0}$ and $\Phi_{p,q}^{\mathbb{Z}^d,1}$, respectively. $\Phi_{p,q}^{\mathbb{Z}^d,0}$ is called the **free**, and $\Phi_{p,q}^{\mathbb{Z}^d,1}$ is called the **wired** random-cluster measure for $\mathbb{Z}^d$ with parameters p and q . These measures are indeed random-cluster measures in the sense of Definition 2.2, moreover, they are extremal among such measures in the following sense.

A natural partial order on the set $\Omega_D = \{0, 1\}^{\mathcal{E}^d}$ of edge configurations is given by defining $\eta' \geq \eta$ for $\eta, \eta' \in \Omega_D$ if for all $e \in \mathcal{E}^d$, $\eta'(e) \geq \eta(e)$. A function $f : \Omega_D \rightarrow \mathbb{R}$ is called **increasing** if $\eta' \geq \eta$ implies $f(\eta') \geq f(\eta)$. For probability measures ϕ, ϕ' on Ω_D , we say that ϕ' is **stochastically larger** than ϕ if for all bounded increasing functions $f : \Omega_D \rightarrow \mathbb{R}$, we have that

$$\int_{\Omega_D} f(\eta) d\phi'(\eta) \geq \int_{\Omega_D} f(\eta) d\phi(\eta).$$

For later purposes we mention that by Strassen's theorem [26] this is equivalent to the existence of an appropriate coupling of the measures ϕ' and ϕ , that is, the existence of a measure Q on $\Omega_D \times \Omega_D$ such that the marginals of Q on the first and second coordinates are ϕ' and ϕ respectively, and $Q(\{(\eta', \eta) \in \Omega_D \times \Omega_D : \eta' \geq \eta\}) = 1$.

It is well-known that $\Phi_{p,q}^{\mathbb{Z}^d,0}$ is the stochastically smallest, and $\Phi_{p,q}^{\mathbb{Z}^d,1}$ is the stochastically largest random-cluster measure for $\mathbb{Z}^d$ with parameters p and q . Therefore, there exists a unique random-cluster measure for $\mathbb{Z}^d$ with parameters p and q if and only if

$$\Phi_{p,q}^{\mathbb{Z}^d,0} = \Phi_{p,q}^{\mathbb{Z}^d,1}. \quad (2)$$

This is the case for any fixed $q \geq 1$, except possibly for at most countably many values of p . It is widely believed that for any $q \geq 1$, there is at most one exceptional p , which can only be the **critical value** $p_c(q, d) = \sup\{p : \Phi_{p,q}^{\mathbb{Z}^d,0}(\{\eta \in \Omega_D : \mathbf{0} \text{ is in an infinite FK cluster in } \eta\}) = 0\}$, where $\mathbf{0}$ denotes the origin in $\mathbb{Z}^d$. It is not difficult to show that the choice of $\Phi_{p,q}^{\mathbb{Z}^d,0}$ in the definition is not crucial. That is, for any random-cluster measure ϕ for $\mathbb{Z}^d$ with parameters p and q , we have that

$$\phi(\{\eta \in \Omega_D : \mathbf{0} \text{ is in an infinite FK cluster in } \eta\}) \begin{cases} = 0 & \text{if } p < p_c(q, d), \\ > 0 & \text{if } p > p_c(q, d). \end{cases}$$

For the rest of the paper, we shall assume without further mentioning that parameters d, p, q for the DaC(q) model on $\mathbb{Z}^d$ are always chosen in such a way that (2) holds, and denote the unique random-cluster measure by $\Phi_{p,q}^{\mathbb{Z}^d}$.

For our main result, we need to consider the critical value in half-spaces as well. Let $\mathcal{H} = \mathcal{H}^d$ denote the subset of $\mathbb{Z}^d$ which consists of the vertices whose first coordinate is strictly positive, and denote $(1, 0, 0, \dots, 0) \in \mathbb{Z}^d$ by u_1 . For $q \geq 1$, we define $p_c^{\mathcal{H}}(q, d) = \sup\{p : \Phi_{p,q}^{\mathbb{Z}^d, 0}(\{\eta \in \Omega_D : u_1 \text{ is in an infinite open path in } \eta \text{ which is contained in } \mathcal{H}\}) = 0\}$. It is obvious that $p_c^{\mathcal{H}}(q, d) \geq p_c(q, d)$. Equality of the two critical values for $q = 1$ was proved by Barsky, Grimmett and Newman [3], for $q = 2$ by Bodineau [4], and for very large values of q , it follows from the Pirogov-Sinai theory (see the last paragraph of Section 2.3 in [4]). For general $q \geq 1$, equality has been conjectured ([25],[19],[4]), but no definite answer is known thus far. However, an upper bound of $p_c^{\mathcal{H}}(q, d) \leq \frac{p_c(1,d)q}{p_c(1,d)q+1-p_c(1,d)}$ can be given easily, using that for $q \geq 1$, $\Phi_{p,q}^{\mathbb{Z}^d, 1}$ is stochastically larger than $\Phi_{\frac{p}{p+(1-p)q}, 1}^{\mathbb{Z}^d, 1}$, and that $p_c^{\mathcal{H}}(1, d) = p_c(1, d)$. Note that $p_c(1, d)$ is the critical value for Bernoulli bond percolation on $\mathbb{Z}^d$. It is well-known (see, e.g., [14]) that for any $d \geq 2$, $0 < p_c(1, d) < 1$. This implies that the above upper bound for $p_c^{\mathcal{H}}(q, d)$ is nontrivial.

Finally, an important feature of the random-cluster measures $\Phi_{p,q}^{\mathbb{Z}^d, 0}$ and $\Phi_{p,q}^{\mathbb{Z}^d, 1}$ with $q \geq 1$ is that they satisfy the FKG inequality [8]. This in particular implies that for $d \geq 2, p \in [0, 1], q \geq 1$, for any edge set $E \subset \mathcal{E}^d$, configuration $\zeta \in \{0, 1\}^E$ on E , and $A_1, A_2 \subset \Omega_D$ increasing events (we call an event $A \subset \Omega_D$ **increasing** if its indicator function is increasing, i.e., if $\eta \in A$ and $\eta' \geq \eta$ implies $\eta' \in A$), we have denoting $B = \{\eta \in \Omega_D : \eta_E = \zeta\}$ that

$$\Phi_{p,q}^{\mathbb{Z}^d, 0}(A_1 \cap A_2 \mid B) \geq \Phi_{p,q}^{\mathbb{Z}^d, 0}(A_1 \mid B) \Phi_{p,q}^{\mathbb{Z}^d, 0}(A_2 \mid B). \quad (3)$$

2.2 Main results

Before stating the main results, let us give the relevant definitions. In this section, μ denotes a probability measure on $\Omega_C = S^{\mathbb{Z}^d}$. Spin configurations will be denoted throughout by ξ, σ , and κ , and the restriction of a spin configuration ξ to a vertex set W by ξ_W . For a set $W \subset \mathbb{Z}^d$ and a spin configuration $\sigma \in S^W$ on W , we denote $K_W^\sigma = \{\xi \in \Omega_C : \xi_W = \sigma\}$. We shall use $A \subset\subset B$ to denote that “ A is a finite subset of B ” throughout. We denote the graph-theoretic distance on $\mathbb{Z}^d$ by dist , and define the **distance** of a vertex $v \in \mathbb{Z}^d$ and a vertex set $H \subset \mathbb{Z}^d$ by $\text{dist}(v, H) = \min\{\text{dist}(v, w) : w \in H\}$. For $k \in \{1, 2, \dots\}$, let $\partial_k H$ denote the k -**neighbourhood** of H , that is, $\partial_k H = \{v \in \mathbb{Z}^d : 1 \leq \text{dist}(H, v) \leq k\}$. Note that $\partial_1 H = \partial H$.

We usually want to view the DaC(q) model as a dependent spin model on $\mathbb{Z}^d$, in which the only role of the edge configuration is to introduce the dependence. One of the first natural questions one may ask about a spin model is whether the finite energy property of [24] holds. This turns out to be the case, moreover, we can even prove a stronger form of it, called uniform nonnullness. The proofs of all statements in this section will be given in Section 4.

Definition 2.3. μ is called **uniformly nonnull** if there exists an $\varepsilon > 0$ such that for all $v \in \mathbb{Z}^d$, $m \in S$, $\sigma \in S^{\mathbb{Z}^d \setminus \{v\}}$, we have

$$\mu(K_{\{v\}}^m \mid K_{\mathbb{Z}^d \setminus \{v\}}^\sigma) \geq \varepsilon.$$

Proposition 2.4. For all $d \in \{1, 2, \dots\}$, $q \geq 1$, $p \in [0, 1)$, and arbitrary values of the other parameters, the measure $\mu_{p,q,(a_1, \dots, a_s)}^{\mathbb{Z}^d}$ is uniformly nonnull.

The concept of k -Markovianity is concerned with the question whether vertices farther than k from a set W have any influence on the spin configuration in W .

Definition 2.5. For $k \in \{1, 2, \dots\}$, μ is called **k -Markovian** if for all $W \subset \subset \mathbb{Z}^d$, $\kappa \in S^W$ and $\sigma, \sigma' \in S^{\mathbb{Z}^d \setminus W}$ such that $\sigma_{\partial_k W} = \sigma'_{\partial_k W}$, we have that

$$\mu(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^\sigma) = \mu(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^{\sigma'}).$$

A weaker notion is that of quasilocality, where the above conditional probabilities do not need to be equal for any k , just their difference is required to tend to 0 as $k \rightarrow \infty$. Due to the compactness of $S^{\mathbb{Z}^d}$ in the product topology, this amounts to the following.

Definition 2.6. μ is called **quasilocal** if for all $W \subset \subset \mathbb{Z}^d$, $\kappa \in S^W$ and $\sigma \in S^{\mathbb{Z}^d \setminus W}$, we have that

$$\lim_{k \rightarrow \infty} \sup_{\substack{\sigma' \in S^{\mathbb{Z}^d \setminus W} \\ \sigma'_{\partial_k W} = \sigma_{\partial_k W}}} |\mu(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^\sigma) - \mu(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^{\sigma'})| = 0.$$

If the above equation holds for μ -almost all $\sigma \in S^{\mathbb{Z}^d \setminus W}$, then μ is called **almost surely quasilocal**.

Finally, we need to say what we mean by Gibbsianness. Instead of the usual definition with absolutely summable interaction potentials (see, e.g., [10, 6]), we shall use a well-known characterisation (see [6], Theorem 2.12),

namely that μ is a **Gibbs measure** if and only if it is quasilocal and uniformly nonnull.

We are now ready to state our main result concerning k -Markovianness and Gibbsianness of the $\text{DaC}(q)$ model. The cases $p = 0, 1$ are trivial, therefore we assume $p \in (0, 1)$. For fixed q, s , and $a_1, \dots, a_s$, recall that $S = \{1, 2, \dots, s\}$, and define $S_{1/q} = \{i \in S : a_i = 1/q\}$. The case $S = S_{1/q}$ is well understood since $S = S_{1/q}$ implies that $s = q$ and $a_1 = a_2 = \dots = a_s$, in which case the procedure defining the $\text{DaC}(q)$ model gives the random-cluster representation of the Potts model. Therefore, for all $p \in (0, 1)$, $\mu_{p,q,(1/q,1/q,\dots,1/q)}^{\mathbb{Z}^d}$ equals a Gibbs measure for the q -state Potts model on $\mathbb{Z}^d$ (at inverse temperature $\beta = -\frac{1}{2} \log(1-p)$). It follows immediately from the standard definition of Potts Gibbs measures with a Hamiltonian (see e.g. [11] for the definition) that all such measures are Markovian (i.e., 1-Markovian). For an alternative proof of the Markovianness of $\mu_{p,q,(1/q,1/q,\dots,1/q)}^{\mathbb{Z}^d}$, see Remark 3.8. If $S \neq S_{1/q}$, let $\ell \in S$ be an (for concreteness, the smallest) index such that $a_\ell = \min\{a_i : i \in S \setminus S_{1/q}\}$.

Theorem 2.7. *Assume that $d \geq 2, q \geq 1$, and that $S \neq S_{1/q}$. Then we have the following.*

1. *For any values of $p, a_1, \dots, a_s \in (0, 1)$, the measure $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is not k -Markovian for any $k \in \mathbb{N}$.*
2. *If $a_\ell > 1/q$, then*
 - (a) *for $p < p_c(qa_\ell, d)$, $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is quasilocal, but*
 - (b) *for $p > p_c^{\mathcal{H}}(qa_\ell, d)$, it is not quasilocal.*
3. *If $a_\ell < 1/q$, then*
 - (a) *if $p < \frac{p_c(1,d)qa_\ell}{p_c(1,d)qa_\ell + 1 - p_c(1,d)}$, then $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is quasilocal, whereas*
 - (b) *if $p > p_c(1, d)$, it is not.*

Combining Theorem 2.7 with Theorem 2.4 and the earlier mentioned characterisation of Gibbs measures, we conclude the following.

Corollary 2.8. *If $S = S_{1/q}$ and in cases 2(a) and 3(a) of Theorem 2.7, the measure $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is a Gibbs measure, and in cases 2(b) and 3(b) of Theorem 2.7, it is not a Gibbs measure.*

The question whether quasilocality is “seriously” violated in cases when $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is not a Gibbs measure (i.e., whether “bad” configurations are exceptional or they actually occur) is related to that of percolation by the following statement, which is a generalisation of Proposition 3.7 in [18].

Proposition 2.9. *Consider the event $E_\infty = \{\xi \in \Omega_C : \xi \text{ contains an infinite connected component of equal spins}\}$. If the parameters $p \in [0, 1]$, $q \geq 1$, $s \in \{2, 3, \dots\}$, and $a_1, \dots, a_s \in (0, 1)$ of the $\text{DaC}(q)$ model are chosen in such a way that*

$$\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(E_\infty) = 0, \quad (4)$$

then $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ satisfies almost sure quasilocality.

It is easy to see that (4) is not a necessary condition for almost sure quasilocality. For instance, one can take $d \geq 2$, $q \geq 1$, $p = 0$, $s = 2$, and an $a_1 < 1$ which is greater than the critical value for Bernoulli *site* percolation on $\mathbb{Z}^d$. Then, although (4) fails, $\mu_{0,q,(a_1,a_2)}^{\mathbb{Z}^d}$ is Markovian (and therefore obviously almost surely quasilocal). Despite this, Proposition 2.9 is not useless. We shall demonstrate this below by giving an application in the two-dimensional case. Häggström's results in Section 3 of [17] imply that for $d = 2$, $q \geq 2$, $p < p_c(q, d)$, if $a_i \leq 1/2$ for all $i \in S$, then (4) holds. Using the main result in [22], this can be extended to $d = 2$, $q \geq 1$, $p < p_c(q, d)$ with the same proof. Combining this with Proposition 2.9, we obtain almost sure quasilocality when $d = 2$ for these parameters.

Corollary 2.10. *If $q \geq 1$, $p < p_c(q, 2)$, and $a_i \leq 1/2$ for all $i \in S$, then $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^2}$ is almost surely quasilocal.*

3 Useful tools

Here we collect the lemmas needed for the proofs of the results in Section 2.2. The statements of the most important ones, Lemma 3.3 and Corollary 3.7, are proved for finite graphs first, then a limit is taken. We shall have an appropriate limiting procedure for $q \geq 1$ only, and this is the reason why we need to restrict to this case in all our results. Throughout this and the next section, we shall use the following notations. For a set $W \subset \mathbb{Z}^d$ and a spin configuration $\sigma \in S^W$ on W , we denote $C_W^\sigma = \{(\xi, \eta) \in \Omega : \xi_W = \sigma\}$. Analogously, for $E \subset \mathcal{E}^d$ and a bond configuration $\zeta \in \{0, 1\}^E$ on E , we denote $D_E^\zeta = \{(\xi, \eta) \in \Omega : \eta_E = \zeta\}$.

For fixed parameters $s \in \{2, 3, \dots\}$, $p, a_1, a_2, \dots, a_s$, and $q \geq 1$, the measure $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ can be obtained as a limit as follows. Let $G_n = (\mathcal{V}_n, \mathcal{E}_n)$ be as in Section 2.1. Consider the $\text{DaC}(q)$ model on G_n with the given parameters as defined in the introduction. Then the corresponding sequence of measures $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{G_n}$ converges to $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ as $n \rightarrow \infty$, in the sense that probabilities of cylinder sets converge. Note that $q \geq 1$ is needed to

ensure the convergence of $\Phi_{p,q}^{G_n}$ to the (unique) random-cluster measure $\Phi_{p,q}^{\mathbb{Z}^d}$, see Section 2.1.

The next two lemmas, which give the conditional edge distribution in the DaC(q) model given any spin configuration, are of crucial importance for the rest of this paper. The statements (and the proofs) are analogues of Proposition 5.1 and Theorem 6.2 in [19]. For a graph $G = (\mathcal{V}, \mathcal{E})$ (where $\mathcal{V}$ and $\mathcal{E}$ are finite or $\mathcal{V} = \mathbb{Z}^d, \mathcal{E} = \mathcal{E}^d$) and a spin configuration $\xi \in \Omega_G^{\mathcal{C}}$ we define for all $i \in S$ the vertex sets $\mathcal{V}^{\xi,i} = \{v \in \mathcal{V} : \xi(v) = i\}$, edge sets $\mathcal{E}^{\xi,i} = \{e = \langle x, y \rangle : x, y \in \mathcal{V}^{\xi,i}\}$ and $\mathcal{E}^{\xi,\text{diff}} = \mathcal{E} \setminus \cup_{i=1}^s \mathcal{E}^{\xi,i}$, and graphs $G^{\xi,i} = (\mathcal{V}^{\xi,i}, \mathcal{E}^{\xi,i})$.

Lemma 3.1. *Let $G = (\mathcal{V}, \mathcal{E})$ be a finite graph. Fix parameters $p \in [0, 1], q > 0, s \in \{2, 3, \dots\}, a_1, a_2, \dots, a_s \in (0, 1)$ in such a way that $\sum_{i=1}^s a_i = 1$, and an arbitrary spin configuration $\sigma \in S^{\mathcal{V}}$, and define the event $A = \{(\xi, \eta) \in \Omega^G : \xi = \sigma\}$. Then we have that*

- (a) *for all $e \in \mathcal{E}^{\sigma,\text{diff}}$, $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^G(\{(\xi, \eta) \in \Omega^G : \eta(e) = 0\} \mid A) = 1$, and*
- (b) *for all $i \in S$, independently for different values of i , on the set $\{0, 1\}^{\mathcal{E}^{\sigma,i}}$, the conditional distribution of $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^G$ given A is the random-cluster measure $\Phi_{p,qa_i}^{G^{\sigma,i}}$.*

Proof. Statement (a) is immediate from the definition of the model. Now let $\eta \in \Omega_D^G$ be such that $\eta(e) = 0$ for all $e \in \mathcal{E}^{\sigma,\text{diff}}$. Denote by $k^{\sigma,i}(\eta)$ the number of connected components in η that have spin i in σ , and notice that $k(\eta) = \sum_{i=1}^s k^{\sigma,i}(\eta)$. Using this observation, (1), and a rearrangement of the factors, we obtain

$$\begin{aligned} \mathbb{P}((\sigma, \eta)) &= \Phi_{p,q}^G(\eta) \prod_{i=1}^s a_i^{k^{\sigma,i}(\eta)} \\ &= \frac{(1-p)^{|\mathcal{E}^{\sigma,\text{diff}}|}}{Z_{p,q}^G} \prod_{i=1}^s \left((qa_i)^{k^{\sigma,i}(\eta)} \prod_{e \in \mathcal{E}^{\sigma,i}} p^{\eta(e)} (1-p)^{1-\eta(e)} \right), \end{aligned}$$

where we wrote $\mathbb{P}$ for $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^G$, and $|\cdot|$ for cardinality. Therefore,

$$\mathbb{P}_{p,q,(a_1,\dots,a_s)}^G((\sigma, \eta) \mid A) = \prod_{i=1}^s \Phi_{p,qa_i}^{G^{\sigma,i}}(\eta_{\mathcal{E}^{\sigma,i}})$$

since the factor $\frac{(1-p)^{|\mathcal{E}^{\sigma,\text{diff}}|} \prod_{i=1}^s Z_{p,q,qa_i}^{G^{\sigma,i}}}{Z_{p,q}^G \mu_{p,q,(a_1,\dots,a_s)}^G(\sigma)}$ is constant in η , thus it must be 1 to give a probability measure. This proves statement (b). $\square$

Remark 3.2. The fact that random-cluster measures factorise on disconnected graphs provides a simple way of drawing a random bond configuration Y with distribution $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^G$ given A . First, set $Y(e) = 0$ for all $e \in \mathcal{E}^{\sigma,\text{diff}}$. Then choose any component $C = (\mathcal{V}_C, \mathcal{E}_C)$ in the graph $(\mathcal{V}, \mathcal{E} \setminus \mathcal{E}^{\sigma,\text{diff}})$. Notice that C is a maximal monochromatic component in G (with respect to σ); suppose that for all $v \in \mathcal{V}_C$, $\sigma(v) = i$. Then, independently of everything else, draw $Y_{\mathcal{E}_C}$ according to the random-cluster measure Φ_{p,qa_i}^C . Repeat this procedure with a new component in $(\mathcal{V}, \mathcal{E} \setminus \mathcal{E}^{\sigma,\text{diff}})$ until there are no more such components. Lemma 3.1 and the observation at the beginning of this paragraph ensure that we get the correct (conditional) distribution.

By using Lemma 3.1 and the limiting procedure for $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$, one obtains analogous statements for $\mathbb{Z}^d$ in case of $q \geq 1$.

Lemma 3.3. *Fix parameters $d, p, q \geq 1, s, (a_1, a_2, \dots, a_s)$ of the DaC(q) model on $\mathbb{Z}^d$ and a spin configuration $\sigma \in \Omega_C$. Then the conditional distribution of $\mathbb{P} = \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $C_{\mathbb{Z}^d}^\sigma$ assigns value 0 to all edges in $\mathcal{E}^{\sigma,\text{diff}}$, and is a random-cluster measure for $G^{\sigma,i}$ with parameters p and qa_i on $\mathcal{E}^{\sigma,i}$, independently for each i . Moreover, for each edge $e \in \mathcal{E}^{\sigma,i}$ and edge configuration $\zeta \in \{0,1\}^{\mathcal{E}^d \setminus \{e\}}$, we have that*

$$\mathbb{P}(\{(\xi, \eta) \in \Omega : \eta(e) = 1\} \mid C_{\mathbb{Z}^d}^\sigma \cap D_{\mathcal{E}^d \setminus \{e\}}^\zeta) = \begin{cases} p & \text{if } x \xleftrightarrow{\zeta} y, \\ \frac{p}{p+(1-p)qa_i} & \text{otherwise,} \end{cases}$$

and the analogue of Remark 3.2 holds in this infinite setting as well.

Proof sketch. Unless the edge configuration $\zeta \in \{0,1\}^{\mathcal{E}^d \setminus \{e\}}$ is special in the sense that it contains at least two infinite FK clusters or there exists an edge $f \in \mathcal{E}^d \setminus \{e\}$ such that changing the state of f in ζ would create at least two infinite FK clusters, we see after a certain stage of the limiting construction described at the beginning of this section whether or not $x \xleftrightarrow{\zeta} y$ occurs, therefore, an equality corresponding to the “moreover” part of Lemma 3.3 can be verified by Lemma 3.1 for all further stages of the limiting construction. Since the above mentioned special edge configurations have $\Phi_{p,q}^{\mathbb{Z}^d}$ -measure 0, we are done. For the details, see the proof of Theorem 6.2 in [19]. $\square$

Lemma 3.3 enables us to give some intuition behind Theorem 2.7. For a vertex set $W \subset \mathbb{Z}^d$, let ΔW denote the **edge boundary** of W , that is, $\Delta W = \{\langle x, y \rangle \in \mathcal{E}^d : x \in W, y \in \mathbb{Z}^d \setminus W\}$. Given a spin configuration σ in $\mathbb{Z}^d \setminus W$, the distribution of spins in W is known once $\sigma_{\partial W}$ and the edge configuration in $W \cup \Delta W$ are known. By Lemma 3.3, the conditional edge distribution in

$\mathcal{E}^d \setminus W$ closes edges between different spin clusters in σ , and is a random-cluster measure with parameters p and qa_i on spin i clusters. If $S = S_{1/q}$, then all such random-cluster measures are product measures with parameter p , so that no spin or edge state outside ∂W matters for the configuration in W . This reasoning will be made precise in Remark 3.8. In any other case, at least one of the underlying random-cluster measures is a dependent bond percolation measure, hence it might be possible to find a configuration σ in which edges far away from W still play a role in determining the states of edges in $W \cup \Delta W$. This will be the case for any p , and for sufficiently large values of p (so large that there is percolation in half-spaces with positive $\Phi_{p, qa_\ell}^{\mathbb{Z}^d, 0}$ -probability if $a_\ell > 1/q$, respectively with positive $\Phi_{p, 1}^{\mathbb{Z}^d, 0}$ -probability if $qa_\ell < 1/q$), we can find edges that play a role at any distance from W in certain configurations.

The next lemma, which is a more general form of Lemma 7.3 in [19], and can be proved in the same way, shows that given edge and spin configurations of a certain type, the “price of changing a spin” depends only on the existence or nonexistence of connections in the edge configuration. Since it looks somewhat specialised, and will not be used until Section 4, the reader might choose to skip it for now.

Lemma 3.4. *Fix parameters $d \geq 2, q \geq 1, p \in [0, 1), s, (a_1, a_2, \dots, a_s)$ of the $DaC(q)$ model, and let $i, j \in S$ be different spins. Then there exist positive constants $c_1^{i,j} = c_1^{i,j}(p, q, a_i, a_j)$ and $c_2^{i,j} = c_2^{i,j}(p, q, a_i, a_j)$ such that for any $v \in \mathbb{Z}^d$ with nearest neighbours $u_1, u_2, \dots, u_{2d}$ and the edges between v and u_i denoted by e_i ($i = 1, 2, \dots, 2d$), we have for all $\sigma \in S^{\mathbb{Z}^d \setminus \{v\}}$ and $\zeta \in \{0, 1\}^{\mathcal{E}^d \setminus \{e_1, e_2, \dots, e_{2d}\}}$ satisfying*

1. $\sigma(u_1) = \sigma(u_2) = i$ and $\sigma(u_3) = \sigma(u_4) = \dots = \sigma(u_{2d}) = j$, and
2. no two of $u_3, u_4, \dots, u_{2d}$ are connected in ζ ,

that

$$\frac{\mathbb{P}_{p, q, (a_1, \dots, a_s)}^{\mathbb{Z}^d}(C_{\{v\}}^i \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma \cap D_{\mathcal{E}^d \setminus \{e_1, \dots, e_{2d}\}}^\zeta)}{\mathbb{P}_{p, q, (a_1, \dots, a_s)}^{\mathbb{Z}^d}(C_{\{v\}}^j \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma \cap D_{\mathcal{E}^d \setminus \{e_1, \dots, e_{2d}\}}^\zeta)} = \begin{cases} c_1^{i,j} & \text{if } u_1 \xleftrightarrow{\zeta} u_2, \\ c_2^{i,j} & \text{otherwise.} \end{cases}$$

The exact values of $c_1^{i,j}$ and $c_2^{i,j}$ are

$$c_1^{i,j} = \frac{p^2 qa_i + 2p(1-p)qa_i + (1-p)^2(qa_i)^2}{(1-p)^2(qa_i)^2} \cdot \frac{a_i}{a_j} \cdot \left(\frac{(1-p)qa_j}{p + (1-p)qa_j} \right)^{2d-2}$$

and

$$c_2^{i,j} = \frac{p^2 + 2p(1-p)qa_i + (1-p)^2(qa_i)^2}{(1-p)^2(qa_i)^2} \cdot \frac{a_i}{a_j} \cdot \left(\frac{(1-p)qa_j}{p + (1-p)qa_j} \right)^{2d-2},$$

and this shows that

$$\begin{cases} c_1^{i,j} > c_2^{i,j} & \text{if and only if } qa_i > 1, \\ c_1^{i,j} = c_2^{i,j} & \text{if and only if } qa_i = 1, \\ c_1^{i,j} < c_2^{i,j} & \text{if and only if } qa_i < 1. \end{cases}$$

Lemma 3.4 will play a role in proving parts 1, 2(b), and 3(b) in Theorem 2.7. For the proof of 2(a) and 3(a), we shall need Lemma 3.10, which is preceded by a few definitions and another lemma. The next definition is motivated by Corollary 3.7.

Definition 3.5. We call an edge set $E = \{e_1, e_2, \dots, e_k\}$ a **barrier** if removing $e_1, e_2, \dots, e_k$ (but not their endvertices) separates the graph $\mathbb{Z}^d$ into two or more disjoint connected subgraphs. Note that exactly one of the resulting subgraphs is infinite, which we call the **exterior** of E , and denote by $\text{ext}(E)$. We denote the vertex set of $\text{ext}(E)$ by $\mathcal{V}_{\text{ext}(E)}$, and the edge set of $\text{ext}(E)$ by $\mathcal{E}_{\text{ext}(E)}$. We call the union of the finite subgraphs the **interior** of E , and denote it by $\text{int}(E)$, and use $\mathcal{V}_{\text{int}(E)}$ and $\mathcal{E}_{\text{int}(E)}$ to denote its vertex and edge set respectively. $E = \{e_1, e_2, \dots, e_k\}$ is called a **closed barrier** in a configuration $(\xi, \eta) \in \Omega$ if E is a barrier and $\eta(e_i) = 0$ holds for $i = 1, \dots, k$, and it is called **quasi-closed barrier** if for all edges $e = \langle x, y \rangle \in E$ such that $\eta(e) = 1$, it is true that $\xi(x) = \xi(y) \in S_{1/q}$.

Note that the edge boundary of a union of finite spin clusters is a closed barrier and that all closed barriers are quasi-closed. According to Lemma 3.3, the states of edges in spin i clusters where $a_i = 1/q$ are chosen independently of everything else, hence they should play no role in issues of dependence. We prove a formal statement concerning this in the following lemma, and we show in Corollary 3.7 a way to make use of this feature of the model. For an event A , we denote the indicator random variable of A by $\mathbb{I}_A$.

Lemma 3.6. Let $G = (\mathcal{V}, \mathcal{E})$ be a finite graph, $V_1, V_2 \subset \mathcal{V}$ a partition of $\mathcal{V}$, and for $i = 1, 2$, define edge sets $E_i = \{e \in \mathcal{E} : \text{both endvertices of } e \text{ are in } V_i\}$, and graphs $G_i = (V_i, E_i)$. Define also the edge set $B = \{e \in \mathcal{E} : e \text{ has one endvertex in } V_1 \text{ and one in } V_2\}$, a subset $B^0 \subset B$, and for $i = 1, 2$, W_i as the set of endvertices of edges in $B \setminus B^0$ that are in V_i . Fix parameters $p, q > 0, s, (a_1, a_2, \dots, a_s)$ of the $\text{DaC}(q)$ model on G , and a spin configuration $\sigma \in S_{1/q}^{W_1 \cup W_2}$ such that for all $e = \langle x, y \rangle \in B \setminus B^0$ we have that $\sigma(x) = \sigma(y)$. Considering the events $C(B, \sigma) = \{(\xi, \eta) \in \Omega^G : \eta_{B^0} \equiv 0, \xi_{W_1 \cup W_2} = \sigma\}$, $K_1 = \{(\xi, \eta) \in \Omega^{G_1} : \xi_{W_1} = \sigma_{W_1}\}$, $K_2 = \{(\xi, \eta) \in \Omega^{G_2} : \xi_{W_2} = \sigma_{W_2}\}$, and

$Z(B^0) = \{\eta \in \{0, 1\}^{B^0} : \eta \equiv 0\}$, we have for each $(\xi, \eta) \in \Omega^G$ that

$$\begin{aligned} \mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^G((\xi, \eta) \mid C(B, \sigma)) &= \mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^{G_1}((\xi_{V_1}, \eta_{E_1}) \mid K_1) \\ &\quad \times \mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^{G_2}((\xi_{V_2}, \eta_{E_2}) \mid K_2) \\ &\quad \times \mathbb{I}_{Z(B^0)}(\eta_{B^0}) \prod_{e \in B \setminus B^0} p^{\eta(e)} (1-p)^{1-\eta(e)}. \end{aligned}$$

This implies in particular the conditional independence given $C(B, \sigma)$ of the random configurations on G_1 and on G_2 .

Proof. Let us fix $(\xi, \eta) \in \Omega^G$. Note that

$$\mathbb{I}_{C(B, \sigma)}(\xi, \eta) = \mathbb{I}_{K_1}(\xi_{V_1}, \eta_{E_1}) \mathbb{I}_{K_2}(\xi_{V_2}, \eta_{E_2}) \mathbb{I}_{Z(B^0)}(\eta_{B^0}),$$

hence if $(\xi, \eta) \notin C(B, \sigma)$ then we have that both sides of the equation that we want to prove are 0, thus for all such configurations we indeed have equality of the two sides. Therefore, let us assume that $(\xi, \eta) \in C(B, \sigma)$. Define the event $A = \{(\kappa, \zeta) \in \Omega^G : \text{there is no edge } e = \langle x, y \rangle \in \mathcal{E} \text{ with } \zeta(e) = 1 \text{ and } \kappa(x) \neq \kappa(y)\}$, and denote the analogously defined subsets of Ω^{G_1} and Ω^{G_2} by A_1 and A_2 respectively. Since $(\xi, \eta) \in C(B, \sigma)$, we have that

$$\mathbb{I}_A(\xi, \eta) = \mathbb{I}_{A_1}(\xi_{V_1}, \eta_{E_1}) \mathbb{I}_{A_2}(\xi_{V_2}, \eta_{E_2}).$$

Therefore, if $(\xi, \eta) \notin A$, we have 0 on both sides of the desired equation in Lemma 3.6 by the definition of the model, so let us assume that $(\xi, \eta) \in A$.

Now denote by n the total number of FK clusters in η , and for all $i \in S$, $j = 1, 2$ the number of FK clusters in η that contain a vertex in V_j with spin i in ξ but no vertex in W_j by n_j^i , and for each $i \in S$ the number of FK clusters in η that contain a vertex in $W_1 \cup W_2$ with spin i in ξ by n_3^i . Throughout this proof, we shall omit the subscripts of the joint measures in the DaC(q) models, e.g., we write $\mathbb{P}^G$ for the measure $\mathbb{P}_{p,q,(a_1,a_2,\dots,a_s)}^G$. Since $(\xi, \eta) \in C(B, \sigma) \cap A$, it immediately follows from the definition of $\mathbb{P}^G$ and the definition (1) of random-cluster measures that

$$\mathbb{P}^G((\xi, \eta)) = \frac{q^n}{Z_{p,q}^G} \left(\prod_{e \in \mathcal{E}} p^{\eta(e)} (1-p)^{1-\eta(e)} \right) \left(\prod_{i=1}^s a_i^{n_1^i + n_2^i + n_3^i} \right). \quad (5)$$

Note that $\mathcal{E} = E_1 \cup E_2 \cup B^0 \cup (B \setminus B^0)$. Since $(\xi, \eta) \in C(B, \sigma) \cap A$, we have that $\prod_{e \in B^0} p^{\eta(e)} (1-p)^{1-\eta(e)} = (1-p)^{|B^0|}$ where $|\cdot|$ denotes cardinality, and $n = \sum_{i=1}^s n_1^i + n_2^i + n_3^i$. Furthermore, it is the case that

$$\prod_{i=1}^s (qa_i)^{n_1^i + n_2^i + n_3^i} = \prod_{i=1}^s (qa_i)^{n_1^i + n_2^i},$$

since for all $i \notin S_{1/q}$ we have $n_3^i = 0$, whereas for all $i \in S_{1/q}$, we have $qa_i = 1$, so the factor $\prod_{i=1}^s (qa_i)^{n_3^i}$ is indeed 1. Using these observations, we can factorise the expression in (5). Indeed, denoting the quantity $(1-p)^{|B^0|}/(Z_{p,q}^G \mathbb{P}^G(C(B, \sigma)))$ which does not depend on (ξ, η) by c , we have that

$$\begin{aligned} \mathbb{P}^G((\xi, \eta) \mid C(B, \sigma)) &= \frac{\mathbb{P}^G((\xi, \eta))}{\mathbb{P}^G(C(B, \sigma))} \\ &= c \left[\prod_{e \in E_1} p^{\eta(e)} (1-p)^{1-\eta(e)} \prod_{i=1}^s (qa_i)^{n_1^i} \right] \\ &\quad \times \left[\prod_{e \in E_2} p^{\eta(e)} (1-p)^{1-\eta(e)} \prod_{i=1}^s (qa_i)^{n_2^i} \right] \\ &\quad \times \left[\prod_{e \in B \setminus B_0} p^{\eta(e)} (1-p)^{1-\eta(e)} \right]. \end{aligned}$$

The last part of the proof, namely to show that the expressions between the first and second pair of square brackets are $c_1 \mathbb{P}^{G_1}((\xi_{V_1}, \eta_{E_1}) \mid K_1)$ and $c_2 \mathbb{P}^{G_2}((\xi_{V_2}, \eta_{E_2}) \mid K_2)$ respectively where c_1 and c_2 are constants (i.e., they do not depend on ξ or η) will be easy. It is sufficient to show the first one, since then the second one follows by relabeling V_1 and V_2 . Let n_4 denote the total number of FK clusters in η_{E_1} , and for each $i \in S$, n_5^i the number of FK clusters in η_{E_1} that contain a vertex in W_1 with spin i in ξ_{V_1} . Since $(\xi, \eta) \in C(B, \sigma) \cap A$, we have that $n_4 = \sum_{i=1}^s n_1^i + n_5^i$. Similarly as in the paragraph after (5), we have $n_5^i = 0$ for all $i \notin S_{1/q}$ and $qa_i = 1$ for all $i \in S_{1/q}$, hence

$$\prod_{i=1}^s (qa_i)^{n_1^i + n_5^i} = \prod_{i=1}^s (qa_i)^{n_1^i}.$$

Denoting $Z_{p,q}^{G_1} \mathbb{P}^{G_1}(K_1)$ by c_1 , the above observations imply that

$$\begin{aligned} \mathbb{P}^{G_1}((\xi_{V_1}, \eta_{E_1}) \mid K_1) &= \frac{\mathbb{P}^{G_1}((\xi_{V_1}, \eta_{E_1}))}{\mathbb{P}^{G_1}(K_1)} \\ &= \frac{q^{n_4}}{c_1} \prod_{e \in E_1} p^{\eta(e)} (1-p)^{1-\eta(e)} \prod_{i=1}^s a_i^{n_1^i + n_5^i} \\ &= \frac{1}{c_1} \prod_{e \in E_1} p^{\eta(e)} (1-p)^{1-\eta(e)} \prod_{i=1}^s (qa_i)^{n_1^i}. \end{aligned}$$

Finally, notice that none of c, c_1, c_2 depends on (ξ, η) , hence the product cc_1c_2 must be equal to 1 to make $\mathbb{P}^G(\cdot \mid C(B, \sigma))$ a probability measure. This observation completes the proof of Lemma 3.6. $\square$

Lemma 3.6 combined with the limiting procedure for $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ yields the following result, which shows why quasi-closed barriers are useful.

Corollary 3.7. *Fix parameters $d, p, q \geq 1, s$, and $(a_1, a_2, \dots, a_s)$ of the $\text{DaC}(q)$ model on $\mathbb{Z}^d$. Let (X, Y) be a random configuration in Ω with distribution $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$, B a barrier, and $C(B)$ the event that B is quasi-closed. Then, given $C(B)$, $(X_{\mathcal{V}_{\text{int}(B)}}, Y_{\mathcal{E}_{\text{int}(B)}})$ and $(X_{\mathcal{V}_{\text{ext}(B)}}, Y_{\mathcal{E}_{\text{ext}(B)}})$ are conditionally independent. In particular, for a set $H \subset \mathbb{Z}^d$ and a spin configuration $\sigma \in S^H$, we have that the conditional distribution of $(X_{\mathcal{V}_{\text{int}(B)}}, Y_{\mathcal{E}_{\text{int}(B)}})$ given $C(B)$ and $\{(\xi, \eta) \in \Omega : \xi_H = \sigma_H\}$ is $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\text{int}(B)}$ conditioned on $\{(\xi, \eta) \in \Omega^{\text{int}(B)} : \xi_{H \cap \mathcal{V}_{\text{int}(B)}} = \sigma_{H \cap \mathcal{V}_{\text{int}(B)}}\}$.*

Remark 3.8. As a first application of Corollary 3.7, we give a proof of the Markovianity of the measure $\mu_{p,q,(1/q,1/q,\dots,1/q)}^{\mathbb{Z}^d}$ that does not use the connection between the $\text{DaC}(q)$ and Potts models. Fix a finite subset W of $\mathbb{Z}^d$. For any spin configuration $\sigma \in S^{\mathbb{Z}^d \setminus W}$ outside W , we have that the edge set $B = \{e \in \mathcal{E}^d : e \text{ has one endvertex in } \partial W \text{ and one in } \partial_2 W\}$ is a quasi-closed barrier since for each edge $e = \langle x, y \rangle \in B$, it is either the case that $\sigma(x) \neq \sigma(y)$ and therefore e is closed, or $\sigma(x) = \sigma(y) \in S_{1/q}$ since $S = S_{1/q}$. Therefore, we have by Corollary 3.7 that for any spin configurations $\sigma, \sigma' \in S^{\mathbb{Z}^d \setminus W}$ with $\sigma'_{\partial W} = \sigma_{\partial W}$, the conditional distributions $\mathbb{P}_{p,q,(1/q,1/q,\dots,1/q)}^{\mathbb{Z}^d}$ given $K_{\mathbb{Z}^d \setminus W}^\sigma$ and $\mathbb{P}_{p,q,(1/q,1/q,\dots,1/q)}^{\mathbb{Z}^d}$ given $K_{\mathbb{Z}^d \setminus W}^{\sigma'}$ are the same in $S^{\mathcal{V}_{\text{int}(B)}} \times \{0, 1\}^{\mathcal{E}_{\text{int}(B)}}$. The statement follows.

Roughly speaking, quasilocality of $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ means that the spin distribution in a set H does not depend on spins very far from H . Corollary 3.7 shows that this is the case if H is surrounded by a quasi-closed barrier. The following definition will help us to localise quasi-closed barriers.

Definition 3.9. *Let (X, Y) be an $S^{\mathbb{Z}^d} \times \{0, 1\}^{\mathcal{E}^d}$ -valued random pair with distribution $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$. Given $(X, Y) = (\xi, \eta)$ for some $(\xi, \eta) \in \Omega$, let $\hat{Y} \in \{0, 1\}^{\mathcal{E}^d}$ be defined by setting, for each $e = \langle x, y \rangle \in \mathcal{E}^d$,*

$$\hat{Y}(e) = \begin{cases} 0 & \text{if } \xi(x) = \xi(y) \in S_{1/q}, \\ \eta(e) & \text{otherwise.} \end{cases}$$

We write $\hat{\mathbb{P}}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ for the induced joint distribution of $(X, Y, \hat{Y})$ on $\hat{\Omega} = S^{\mathbb{Z}^d} \times \{0, 1\}^{\mathcal{E}^d} \times \{0, 1\}^{\mathcal{E}^d}$.

Now recall the definition of $\ell \in S$ right before 2.7 and of C_W^σ and D_E^ζ at the beginning of Section 3. The next lemma, which is a generalisation of Lemma 9.5 in [19], compares the conditional distribution of $\hat{Y}$ given a spin

configuration and $\hat{Y}$ outside a finite edge set F to a random-cluster measure for $\mathbb{Z}^d$ with parameters p and qa_ℓ in case of $a_\ell > 1/q$, and the conditional distribution of Y given a spin configuration and Y outside a finite edge set F to a Bernoulli bond percolation measure with parameter $\frac{p}{p+(1-p)qa_\ell}$ if $a_\ell < 1/q$.

Lemma 3.10. *Suppose that $q \geq 1$ and $S \neq S_{1/q}$. For any spin configuration $\sigma \in S^{\mathbb{Z}^d}$, edge set $F \subset \mathcal{E}^d$, and edge configurations $\zeta, \zeta' \in \{0, 1\}^{\mathcal{E}^d \setminus F}$ such that $\zeta' \geq \zeta$, denoting $\hat{A} = \{(\xi, \eta, \hat{\eta}) \in \hat{\Omega} : \xi = \sigma, \hat{\eta}_{\mathcal{E}^d \setminus F} = \zeta\}$ and $A' = \{\eta \in \Omega_D : \eta_{\mathcal{E}^d \setminus F} = \zeta'\}$, we have the following.*

1. *If $a_\ell > 1/q$, let ϕ be a random-cluster measure for $\mathbb{Z}^d$ with parameters p and qa_ℓ . Then the conditional distribution of ϕ given A' is stochastically larger than the marginal on $\hat{Y}$ of $\hat{\mathbb{P}}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $\hat{A}$.*
2. *If $a_\ell < 1/q$, then the conditional distribution of the product measure $\Phi_{\frac{p}{p+(1-p)qa_\ell}, 1}^{\mathbb{Z}^d, 1}$ given A' is stochastically larger than the marginal on Y of the conditional distribution of $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $C_{\mathbb{Z}^d}^\sigma \cap D_{\mathcal{E}^d \setminus F}^\zeta$.*

Proof. First we prove part 1. By Holley's theorem (see [11], Theorem 4.8), it is sufficient to prove that for all $a \in \{0, 1\}$, $e = \langle x, y \rangle \in F$, $\zeta_g, \zeta_s \in \{0, 1\}^{F \setminus \{e\}}$ such that $\zeta_g \geq \zeta_s$, denoting $B_g = \{\eta \in \Omega_D : \eta_{F \setminus \{e\}} = \zeta_g\}$ and $\hat{B}_s = \{(\xi, \eta, \hat{\eta}) \in \hat{\Omega} : \hat{\eta}_{F \setminus \{e\}} = \zeta_s\}$, we have that

$$\phi(\{\eta \in \Omega_D : \eta(e) \geq a\} \mid A' \cap B_g) \quad (6)$$

is greater than or equal to

$$\hat{\mathbb{P}}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(\{(\xi, \eta, \hat{\eta}) \in \hat{\Omega} : \hat{\eta}(e) \geq a\} \mid \hat{A} \cap \hat{B}_s). \quad (7)$$

This is obvious for $a = 0$. For $a = 1$, using the notation $(\eta_1; \eta_2)$ for an edge configuration which agrees with η_1 on $\mathcal{E}^d \setminus F$ and with η_2 on $F \setminus \{e\}$, we have by Definition 2.2 of random-cluster measures that (6) equals

$$\begin{cases} p & \text{if } x \xleftrightarrow{(\zeta'; \zeta_g)} y, \\ \frac{p}{p+(1-p)qa_\ell} & \text{if } x \xleftrightarrow{(\zeta'; \zeta_g)} y. \end{cases}$$

For (7), we need to check first what the spins of the endvertices x, y of e are in σ . Indeed, if $\sigma(x) \neq \sigma(y)$ or $\sigma(x) = \sigma(y) \in S_{1/q}$, then (7) = 0 by Definition 3.9. Let us assume that $\sigma(x) = \sigma(y) = j \notin S_{1/q}$, and denote the maximal monochromatic component (with respect to σ) in the graph $\mathbb{Z}^d$ which contains x by G_x . By Lemma 3.3, the conditional distribution of

Y given σ is a random-cluster measure on G_x with parameters p and qa_j . Moreover, since $j \notin S_{1/q}$, we have that $\hat{Y}$ and Y agree on G_x . Keeping these observations in mind, it follows that (7) equals

$$\begin{cases} 0 & \text{if } \sigma(x) \neq \sigma(y) \text{ or } \sigma(x) = \sigma(y) \in S_{1/q}, \\ p & \text{if } \sigma(x) = \sigma(y) \notin S_{1/q} \text{ and } x \stackrel{(\zeta; \zeta_s)}{\longleftrightarrow} y, \\ \frac{p}{p+(1-p)qa_j} & \text{if } \sigma(x) = \sigma(y) = j \notin S_{1/q} \text{ and } x \stackrel{(\zeta; \zeta_s)}{\nleftrightarrow} y. \end{cases}$$

Since due to the assumption $a_\ell \geq 1/q$, we have for all $j \in S$ that

$$p \geq \frac{p}{p+(1-p)qa_j}$$

and by the definition of a_ℓ we have for all $j \in S \setminus S_{1/q}$ that

$$\frac{p}{p+(1-p)qa_\ell} \geq \frac{p}{p+(1-p)qa_j},$$

we obtain the desired result by noting that $x \stackrel{(\zeta'; \zeta_g)}{\nleftrightarrow} y$ implies $x \stackrel{(\zeta; \zeta_s)}{\nleftrightarrow} y$.

Part 2 can also be proved by a direct application of Holley's theorem, noticing that due to the definition of ℓ and the assumption $qa_\ell < 1$ we have that

$$\frac{p}{p+(1-p)qa_\ell} \geq \max\{p, \max_{i \in S} \frac{p}{p+(1-p)qa_i}\}.$$

□

4 Proofs of the main results

After all the preparation in Section 3, we are now ready to prove our main results. The proof of Proposition 2.4 is not difficult. In fact, one can use the same idea as the one behind the proof of Lemma 5.6 in [18], namely that any vertex can be isolated (i.e., incident to closed edges only) in the edge configuration (given any spin configuration) with probability bounded away from 0, in which case it can be assigned any spin in S , independently of everything else. A formal proof goes as follows.

Proof of Proposition 2.4. Fix $v \in \mathbb{Z}^d, m \in S$ and $\sigma \in S^{\mathbb{Z}^d \setminus \{v\}}$, and recall the definition for $W \subset \mathbb{Z}^d$ of the event $K_W^\sigma \subset \Omega_C$ at the beginning of Section 2.2 and of the analogous event $C_W^\sigma \subset \Omega$ at the beginning of Section 3. Denote by E_v the event that all $2d$ edges incident to v are closed. We have that

$$\begin{aligned} \mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_{\{v\}}^m \mid K_{\mathbb{Z}^d \setminus \{v\}}^\sigma) &\geq \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(C_{\{v\}}^m \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma \cap E_v) \\ &\quad \times \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(E_v \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma). \end{aligned} \quad (8)$$

Obviously (or as a special case of Corollary 3.7), we have that the first term on the right hand side of (8) is a_m , since given E_v , v is assigned a spin independently of everything else.

On the other hand,

$$\begin{aligned} \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(E_v \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma) &= \sum_{b \in S} \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(E_v \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma \cap C_{\{v\}}^b) \\ &\quad \times \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(C_{\{v\}}^b \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma). \end{aligned}$$

Now, whatever value $b \in S$ takes, the full spin configuration is given in the first factor on the right hand side, so we can apply Lemma 3.3. Under any random-cluster measure with parameters p and $\tilde{q} > 0$, the probability of E_v is bounded away from 0: a lower bound for $\tilde{q} \geq 1$ is $(1-p)^{2d}$, while for $\tilde{q} < 1$ it is $(1 - \frac{p}{p+(1-p)\tilde{q}})^{2d}$. Since here the parameter $\tilde{q}$ equals qa_b for some b , we get the lower bound

$$\min\{(1-p)^{2d}, (1 - \frac{p}{p + (1-p)q \min_{i \in S} a_i})^{2d}\}$$

for the first factor, which is uniform in b . As $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(C_{\{v\}}^b \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma)$ for $b \in S$ sum up to 1, we get the same lower bound for $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(E_v \mid C_{\mathbb{Z}^d \setminus \{v\}}^\sigma)$.

Combining this with (8) and the remark thereafter, we have that

$$\varepsilon = \left(\min_{i \in S} a_i \right) \left(\min\{1-p, 1 - \frac{p}{p + (1-p)q \min_{i \in S} a_i}\} \right)^{2d}$$

is a lower bound for $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_{\{v\}}^m \mid K_{\mathbb{Z}^d \setminus \{v\}}^\sigma)$. Since ε does not depend on v, m or σ and is positive for any values of $p \in [0, 1], q \geq 1, a_1, \dots, a_s \in (0, 1)$, we conclude that $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is uniformly nonnull for such parameters. $\square$

The proof of Theorem 2.7 consists of many parts. For the proof of parts 1, 2(b), and 3(b), we use a counterexample that is very similar to the one given in [18, 19] (see also [7, 20, 21]), defined below. After the definition, we give Lemma 4.1 after which it will not be difficult to prove parts 1, 2(b), and 3(b). Finally, we prove parts 2(a) and 3(a). From this point on we assume that $S \neq S_{1/q}$.

Recall first the definitions of Λ_n and ℓ from Sections 2.1 and 2.2, respectively. Fix an arbitrary spin $m \in S$ such that $m \neq \ell$, and define an auxiliary spin configuration $\sigma^* \in S^{\mathbb{Z}^d}$ by setting, for each $x = (x_1, x_2, \dots, x_d) \in \mathbb{Z}^d$,

$$\sigma^*(x) = \begin{cases} m & \text{if } x_1 = 0, |x_2| + |x_3| + \dots + |x_d| = 1 \\ & \text{or } x_1 = -1, |x_2| + |x_3| + \dots + |x_d| > 1, \text{ and} \\ \ell & \text{otherwise,} \end{cases}$$

and for $k = 1, 2, \dots$, spin configurations $\sigma^{k,\ell}, \sigma^{k,m} \in S^{\mathbb{Z}^d \setminus \{\mathbf{0}\}}$ (see Figure 1 below) by

$$\sigma^{k,\ell}(x) = \begin{cases} \ell & \text{for } x \in \mathbb{Z}^d \setminus \Lambda_k, \\ \sigma^*(x) & \text{otherwise,} \end{cases}$$

and

$$\sigma^{k,m}(x) = \begin{cases} m & \text{for } x \in \Lambda_{k+1} \setminus \Lambda_k, \\ \sigma^{k,\ell}(x) & \text{otherwise.} \end{cases}$$




Figure 1: Restriction of $\sigma^{3,\ell}$ (to the left) and $\sigma^{3,m}$ (to the right) to $\Lambda_4 \setminus \{\mathbf{0}\}$ in two dimensions. For all $x \in \mathbb{Z}^2 \setminus \Lambda_4$, $\sigma^{3,\ell}(x) = \sigma^{3,m}(x) = \ell$.

Denote the two nearest neighbours of $\mathbf{0}$ in $\mathbb{Z}^d$ with ξ^* -spin ℓ by $u_1 = (1, 0, 0, \dots, 0)$ and $u_2 = (-1, 0, 0, \dots, 0)$, the other nearest neighbours by $u_3, u_4, \dots, u_{2d}$, and for $i = 1, 2, \dots, 2d$ the edges between $\mathbf{0}$ and u_i by e_i . Most of the work needed for the proof of parts 1, 2(b), and 3(b) of Theorem 2.7 is contained in the following lemma.

Lemma 4.1. *Fix parameters $d \geq 2, q \geq 1, p, a_1, a_2, \dots, a_s \in (0, 1)$ of the $\text{DaC}(q)$ model on $\mathbb{Z}^d$ in such a way that $S \neq S_{1/q}$. Considering the events $A = \{(\xi, \eta) \in \Omega : \text{there exists an open path in } \eta_{\mathbb{Z}^d \setminus \{e_1, e_2, \dots, e_{2d}\}} \text{ between } u_1 \text{ and } u_2\}$ and $O^{\ell,m} = \{(\xi, \eta) \in \Omega : \xi(\mathbf{0}) \in \{\ell, m\}\}$, we have the following.*

1. *If for a fixed $k \in \{1, 2, \dots\}$ we have*

$$\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(A \mid O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}) > 0,$$

then $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is not k -Markovian.

2. *If there exists $\gamma > 0$ such that, for all $k \in \{1, 2, \dots\}$,*

$$\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(A \mid O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}) > \gamma,$$

then $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is not quasilocal.

Proof. In order to simplify the notation, in this proof we denote $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ by $\mathbb{P}$, $C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}$ by $L = L^k$, and $C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,m}}$ by $M = M^k$. The first step in the proof is to derive, for all $k \in \{1, 2, \dots\}$, inequality (10). Consider the expression

$$|\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | M)|. \quad (9)$$

Note that we have

$$\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) = \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L) \mathbb{P}(O^{\ell,m} | L),$$

and similarly with M . Using this, we obtain by basic algebra (i.e., first subtracting, then adding a dummy term $\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M) \mathbb{P}(O^{\ell,m} | L)$ in (9) between the absolute values and finally using that $|a - b| \geq |a| - |b|$) that (9) is greater than or equal to

$$\begin{aligned} & |\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M)| \mathbb{P}(O^{\ell,m} | L) \\ & - |\mathbb{P}(O^{\ell,m} | L) - \mathbb{P}(O^{\ell,m} | M)| \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M). \end{aligned}$$

Since $\mathbb{P}(O^{\ell,m} | L) = \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) + \mathbb{P}(C_{\{\mathbf{0}\}}^m | L)$, we have by uniform nonnullness (i.e., Proposition 2.4) that there exists $\delta > 0$ such that, uniformly in k , $|\mathbb{P}(O^{\ell,m} | L)| \geq \delta$. This observation, noting that $\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M) \leq 1$, and that $\mathbb{P}(O^{\ell,m} | L) = \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) + \mathbb{P}(C_{\{\mathbf{0}\}}^m | L)$ and similarly with M , and applying the triangle inequality yields that (9) is greater than or equal to

$$\begin{aligned} & |\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M)| \delta \\ & - |\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | M)| - |\mathbb{P}(C_{\{\mathbf{0}\}}^m | L) - \mathbb{P}(C_{\{\mathbf{0}\}}^m | M)|. \end{aligned}$$

After a rearrangement of the terms, this gives that

$$\begin{aligned} & 2 |\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | M)| + |\mathbb{P}(C_{\{\mathbf{0}\}}^m | L) - \mathbb{P}(C_{\{\mathbf{0}\}}^m | M)| \\ & \geq \delta |\mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap M)|. \end{aligned} \quad (10)$$

From now on, we shall be working on bounding the right hand side of (10) from below. Elementary calculations and an application of Lemma 3.4 with $i = \ell, j = m, v = \mathbf{0}$ show that

$$\begin{aligned} \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L \cap A) &= \frac{\mathbb{P}(C_{\{\mathbf{0}\}}^\ell \cap L \cap A) / \mathbb{P}(O^{\ell,m} \cap L \cap A)}{\mathbb{P}(C_{\{\mathbf{0}\}}^m \cap L \cap A) / \mathbb{P}(O^{\ell,m} \cap L \cap A)} \\ &\quad \times \mathbb{P}(C_{\{\mathbf{0}\}}^m | O^{\ell,m} \cap L \cap A) \\ &= c_1^{\ell,m} (1 - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell | O^{\ell,m} \cap L \cap A)), \end{aligned}$$

and therefore

$$\mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid O^{\ell,m} \cap L \cap A) = \frac{c_1^{\ell,m}}{c_1^{\ell,m} + 1}. \quad (11)$$

By similar considerations, we obtain

$$\mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid O^{\ell,m} \cap L \cap A^c) = \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1}, \quad (12)$$

and

$$\begin{aligned} \mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid O^{\ell,m} \cap M) &= \mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid O^{\ell,m} \cap M \cap A^c) \\ &= \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1}. \end{aligned} \quad (13)$$

Using (11) and (12), we get that

$$\begin{aligned} \mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid O^{\ell,m} \cap L) &= \frac{c_1^{\ell,m}}{c_1^{\ell,m} + 1} \mathbb{P}(A \mid O^{\ell,m} \cap L) \\ &\quad + \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1} \mathbb{P}(A^c \mid O^{\ell,m} \cap L) \\ &= \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1} + \left(\frac{c_1^{\ell,m}}{c_1^{\ell,m} + 1} - \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1} \right) \\ &\quad \times \mathbb{P}(A \mid O^{\ell,m} \cap L). \end{aligned} \quad (14)$$

Applying (14) and (13) in (10) yields that, for any k , we have

$$\begin{aligned} &2 \left| \mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid L^k) - \mathbb{P}(C_{\{\mathbf{0}\}}^\ell \mid M^k) \right| + \left| \mathbb{P}(C_{\{\mathbf{0}\}}^m \mid L^k) - \mathbb{P}(C_{\{\mathbf{0}\}}^m \mid M^k) \right| \\ &\geq \delta \left| \frac{c_1^{\ell,m}}{c_1^{\ell,m} + 1} - \frac{c_2^{\ell,m}}{c_2^{\ell,m} + 1} \right| \mathbb{P}(A \mid O^{\ell,m} \cap L^k). \end{aligned} \quad (15)$$

Since $a_\ell \neq 1/q$ by definition, we have that $c_1^{\ell,m} \neq c_2^{\ell,m}$. This implies that the first two factors on the right hand side of (15) are positive constants, neither of which depends on k . Now suppose that $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is k -Markovian for some k . In that case the left hand side of (15) is 0 since $\sigma_{\Lambda_k \setminus \{\mathbf{0}\}}^{k,\ell} = \sigma_{\Lambda_k \setminus \{\mathbf{0}\}}^{k,m}$, therefore $\mathbb{P}(A \mid O^{\ell,m} \cap L^k) = 0$. This proves part 1 of Lemma 4.1. Similarly, if $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is quasilocal, then the limit of the left hand side of (15) is 0 as $k \rightarrow \infty$, which cannot be the case if $\mathbb{P}(A \mid O^{\ell,m} \cap L^k)$ is bounded away from 0, uniformly in k . This concludes the proof of part 2. $\square$

Proof of Theorem 2.7, parts 1, 2(b), and 3(b). For this proof, recall the notion of an increasing event on Ω_D (see Section 2.1). Let $d \geq 2, q \geq 1, p, a_1, a_2, \dots, a_s \in (0, 1)$ be arbitrary parameters of the DaC(q) model on $\mathbb{Z}^d$ in such a way that $S \neq S_{1/q}$. For $k = 1, 2, \dots$, define the edge sets $E^{d,k} = \{e \in \mathcal{E}^d : e \text{ is incident to } \mathbf{0} \text{ or to some } v \in \Lambda_k \setminus \{\mathbf{0}\} \text{ with } \sigma^*(v) = m\}$, with σ^* as defined before Lemma 4.1. For a parameter $\tilde{p} \in (0, 1)$ and each $k = 1, 2, \dots$ we define an inhomogeneous bond percolation measure $P_{\tilde{p},k}$ on Ω_D which assigns value 0 to all $e \in E^{d,k}$, and independently to each $e \in \mathcal{E}^d \setminus E^{d,k}$ value 1 with probability $\tilde{p}$ and 0 with probability $1 - \tilde{p}$. It follows from Lemma 3.3 and Definition 2.2 that the marginal on Y of the conditional distribution $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}$ is stochastically larger than $P_{\tilde{p},k}$ with $\tilde{p} = \frac{p}{p+(1-p)qa_\ell}$ if $qa_\ell \geq 1$, and with $\tilde{p} = p$ if $qa_\ell \leq 1$. Therefore, denoting the projection of A on Ω_D by A_D (note that $A_D \subset \Omega_D$ is increasing), we have for any k that

$$\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(A \mid O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}) \geq P_{\tilde{p},k}(A_D). \quad (16)$$

Since $\min\{\frac{p}{p+(1-p)qa_\ell}, p\} > 0$ for all $p \in (0, 1)$, we obviously have for any fixed $k = 1, 2, \dots$ that $P_{\tilde{p},k}(A_D) > 0$. This proves non- k -Markovianess of the measure $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ according to (16) and part 1 of Lemma 4.1.

For the proof of part 2(b), recall the definition of the vertices $u_1, u_2 \in \mathbb{Z}^d$ and edges $e_1, e_2, \dots, e_{2d} \in \mathcal{E}^d$ (right before Lemma 4.1), and that $\mathcal{H}$ denotes the set of vertices in $\mathbb{Z}^d$ whose first coordinate is strictly positive. Define $\mathcal{H}^-$ as the set of vertices in $\mathbb{Z}^d$ whose first coordinate is strictly negative. Consider the following events:

$$\begin{aligned} A_{\mathcal{H}} &= \{\eta \in \Omega_D : \exists \text{ an infinite open path in } \eta_{\mathcal{H}} \text{ which contains } u_1\}, \\ A_{\mathcal{H}^-} &= \{\eta \in \Omega_D : \exists \text{ an infinite open path in } \eta_{\mathcal{H}^-} \text{ which contains } u_2\}, \\ U &= \{\eta \in \Omega_D : \text{there is at most one infinite open cluster in } \eta_{\mathcal{E}^d \setminus \{e_1, e_2, \dots, e_{2d}\}}\}, \end{aligned}$$

and note that $A_{\mathcal{H}} \cap A_{\mathcal{H}^-} \cap U \subset A_D$.

Now assume that $a_\ell > 1/q$. This implies that $qa_\ell \geq 1$ and hence the free random-cluster measure $\Phi_{p,qa_\ell}^{\mathbb{Z}^d,0}$ exists and is the stochastically smallest random-cluster measure for $\mathbb{Z}^d$ with parameters p and qa_ℓ (see Section 2.1). Let us denote the measure $\Phi_{p,qa_\ell}^{\mathbb{Z}^d,0}$ conditioned on the event $\{\eta \in \Omega_D : \eta_{E^{d,k}} \equiv 0\}$ by $\Phi^{(c)}$. Due to the above mentioned extremality of $\Phi_{p,qa_\ell}^{\mathbb{Z}^d,0}$ with respect to stochastic ordering, Lemma 3.3 implies that the marginal on Y of the measure $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ conditioned on $O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}$ is stochastically larger than $\Phi^{(c)}$. Therefore, we have that

$$\begin{aligned} \mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(A \mid O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{k,\ell}}) &\geq \Phi^{(c)}(A_D) \\ &\geq \Phi^{(c)}(A_{\mathcal{H}} \cap A_{\mathcal{H}^-} \cap U). \end{aligned} \quad (17)$$

It is not difficult to see that the fact that under $\Phi_{p,qa_\ell}^{\mathbb{Z}^d,0}$ there is a.s. at most one infinite FK cluster and closing all the edges $e_1, e_2, \dots, e_{2d}$ cannot change this implies the same thing for $\Phi^{(c)}$. Hence, we have that

$$\begin{aligned}\Phi^{(c)}(A_{\mathcal{H}} \cap A_{\mathcal{H}^-} \cap U) &= \Phi^{(c)}(A_{\mathcal{H}} \cap A_{\mathcal{H}^-}) \\ &\geq \Phi^{(c)}(A_{\mathcal{H}}) \Phi^{(c)}(A_{\mathcal{H}^-})\end{aligned}\tag{18}$$

by (3), since $A_{\mathcal{H}}$ and $A_{\mathcal{H}^-}$ are increasing events. Now note that it follows from the definition of $p_c^{\mathcal{H}}(qa_\ell, d)$ that for all $p > p_c^{\mathcal{H}}(qa_\ell, d)$, $\Phi^{(c)}(A_{\mathcal{H}}) = \Phi^{(c)}(A_{\mathcal{H}^-})$ is bounded away from 0, uniformly in k . Therefore, by (17) and (18), $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(A \mid O^{\ell,m} \cap C_{\mathbb{Z}^d \setminus \{\mathbf{0}\}}^{\sigma^{\kappa,\ell}})$ is bounded away from 0 for such values of p , which implies non-quasilocality of $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ by part 2 of Lemma 4.1. This concludes the proof of part 2(b).

In case of $a_\ell < 1/q$, as remarked above, (16) holds with $\tilde{p} = p$. On the other hand, if $p > p_c(1, d)$, then $\tilde{p} > p_c(1, d)$, hence by Lemma 8.2 in [19] (whose proof is based on a computation similar to (17) and (18)), we have

$$\lim_{k \rightarrow \infty} P_{\tilde{p},k}(A_D) > 0.$$

By this, (16), and part 2 of Lemma 4.1, it follows that $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ is not quasilocal, proving part 3(b). $\square$

Proof of Theorem 2.7, parts 2(a) and 3(a). For the proof, recall the definition of $\hat{\Omega}$ (Definition 3.9), and for a set $W \subset \mathbb{Z}^d$ and a spin configuration $\kappa \in S^W$, the definition of K_W^κ (Section 2.2), and define the analogous event $\hat{C}_W^\kappa = \{(\xi, \eta, \hat{\eta}) \in \hat{\Omega} : \xi_W = \kappa\}$. Fix parameters $d \geq 2, q \geq 1, a_1, a_2, \dots, a_s \in (0, 1)$ of the DaC(q) model on $\mathbb{Z}^d$, then p in such a way that $p < p_c(qa_\ell, d)$ if $a_\ell > 1/q$, and $p < \frac{p_c(1,d)qa_\ell}{p_c(1,d)qa_\ell + 1 - p_c(1,d)}$ if $a_\ell < 1/q$. Fix an arbitrary $W \subset \subset \mathbb{Z}^d$, $\kappa \in S^W$, and $\varepsilon > 0$. We need to show the existence of $N = N(\varepsilon, W, \kappa)$ such that for all $n \geq N$, if $\sigma, \sigma' \in S^{\mathbb{Z}^d \setminus W}$ agree on $\Lambda_n \setminus W$, then

$$|\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^\sigma) - \mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^{\sigma'})| \tag{19}$$

is smaller than or equal to ε .

In order to find such an N , following the idea in the proof of Theorem 4.4 part (i) in [19], we shall consider a measure on Ω_D that is stochastically larger than the conditional distribution of edges given either of σ or σ' : we define $\phi^{\text{dom}} = \Phi_{p,qa_\ell}^{\mathbb{Z}^d,1}$ in case of $a_\ell > 1/q$, and $\phi^{\text{dom}} = \Phi_{\frac{p}{p+(1-p)qa_\ell},1}^{\mathbb{Z}^d,1}$ if $a_\ell < 1/q$. Note that the parameters are chosen in such a way that ϕ^{dom} -a.s. there exists no infinite open bond cluster (for the case $a_\ell < 1/q$, notice that $p <$

$\frac{p_c(1,d)qa_\ell}{p_c(1,d)qa_\ell+1-p_c(1,d)}$ ensures that $\frac{p}{p+(1-p)qa_\ell} < p_c(1,d)$. Therefore, it is possible to choose an N so large that

$$\phi^{\text{dom}}(\{\partial W \leftrightarrow \partial \Lambda_N\}) \leq \varepsilon, \quad (20)$$

where $\{\partial W \leftrightarrow \partial \Lambda_N\} = \{\eta \in \Omega_D : \text{there exists a path between } \partial W \text{ and } \partial \Lambda_N \text{ along which all edges are open in } \eta\}$.

Fix an arbitrary $n \geq N$, and let $\sigma, \sigma' \in S^{\mathbb{Z}^d \setminus W}$ be two arbitrary spin configurations such that $\sigma_{\Lambda_n \setminus W} = \sigma'_{\Lambda_n \setminus W}$. We construct a coupling of

- (i) an $\hat{\Omega}$ -valued random triple $(X, Y, \hat{Y})$ with distribution $\hat{\mathbb{P}}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ conditioned on $\hat{C}_{\mathbb{Z}^d \setminus W}^\sigma$,
- (ii) an $\hat{\Omega}$ -valued random triple $(X', Y', \hat{Y}')$ with distribution $\hat{\mathbb{P}}_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ conditioned on $\hat{C}_{\mathbb{Z}^d \setminus W}^{\sigma'}$,
- (iii) an Ω_D -valued random edge configuration Y^{dom} with distribution ϕ^{dom} ,

and write $\mathbb{Q}$ for the probability measure on $\hat{\Omega} \times \hat{\Omega} \times \Omega_D$ underlying the coupling. It follows from the coupling inequality (Proposition 4.2 in [11]) that (19) $\leq \mathbb{Q}(X_W \neq X'_W)$, hence showing that $\mathbb{Q}(X_W = X'_W) \geq 1 - \varepsilon$ would complete the proof. The idea is the following.

1. Draw $\hat{Y}, \hat{Y}'$ and Y^{dom} outside Λ_n in such a way that

$$\mathbb{Q}(\hat{Y}_{\mathcal{E}^d \setminus \mathcal{E}_n} \leq Y_{\mathcal{E}^d \setminus \mathcal{E}_n}^{\text{dom}} \text{ and } \hat{Y}'_{\mathcal{E}^d \setminus \mathcal{E}_n} \leq Y_{\mathcal{E}^d \setminus \mathcal{E}_n}^{\text{dom}}) = 1.$$

This is possible due to Lemma 3.10 part 1 if $qa_\ell > 1$, part 2 if $qa_\ell < 1$, and Strassen's theorem (see Section 2.1).

2. Determine the statuses of the edges inside the box, one edge at a time, in a random order determined by Y^{dom} : at each step, we choose the first edge e (according to a deterministic ordering of $\mathcal{E}_n$) which has not been selected in any previous step and is incident to some vertex in $\partial \Lambda_n$ or some previously selected edge f with $Y^{\text{dom}}(f) = 1$. This roughly means that the statuses of edges are checked going from $\partial \Lambda_n$ towards the centre of the box until a barrier is found which is closed in Y^{dom} or there are no more edges left. The event T that there exists a barrier B that separates ∂W and $\partial \Lambda_n$ (i.e., such that $W \cup \partial W \subset \mathcal{V}_{\text{int}(B)} \subset \Lambda_n$) which is closed in Y^{dom} occurs with probability at least $1 - \varepsilon$ by (20), so let us assume that. Using Lemma 3.10 part 1 if $qa_\ell > 1$ and part 2 if $qa_\ell < 1$ at each step, it is possible to ensure that $\hat{Y}(e) \leq Y^{\text{dom}}(e)$ and $\hat{Y}'(e) \leq Y^{\text{dom}}(e)$ for each edge e selected in this stage. This in particular means that B is closed in $\hat{Y}$ and $\hat{Y}'$ as well, i.e., for all $e \in B$, $\hat{Y}(e) = \hat{Y}'(e) = 0$.

3. Since B is a barrier which closed in $\hat{Y}$, it is a quasi-closed barrier in (X, Y) , therefore Corollary Lemma 3.7 implies that the conditional distribution of (X, Y) on $\text{int}(B)$ given $X_{\mathbb{Z}^d \setminus W} = \sigma$ and what we have seen of $(X, Y, \hat{Y})$ is $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\text{int}(B)}$ conditioned on $\{(\xi, \eta) \in \Omega^{\text{int}(B)} : \xi_{\mathcal{V}_{\text{int}(B)} \setminus W} = \sigma_{\mathcal{V}_{\text{int}(B)} \setminus W}\}$. By similar considerations, the conditional distribution of (X', Y') given what we have seen of $(X', Y', \hat{Y}')$ is $\mathbb{P}_{p,q,(a_1,\dots,a_s)}^{\text{int}(B)}$ conditioned on $\{(\xi, \eta) \in \Omega^{\text{int}(B)} : \xi_{\mathcal{V}_{\text{int}(B)} \setminus W} = \sigma'_{\mathcal{V}_{\text{int}(B)} \setminus W}\}$. Since $\mathcal{V}_{\text{int}(B)} \subset \Lambda_n$ and $\sigma_{\Lambda_n \setminus W} = \sigma'_{\Lambda_n \setminus W}$, we can take

$$(X_{\mathcal{V}_{\text{int}(B)}}, Y_{\mathcal{E}_{\text{int}(B)}}, \hat{Y}_{\mathcal{E}_{\text{int}(B)}}) = (X'_{\mathcal{V}_{\text{int}(B)}}, Y'_{\mathcal{E}_{\text{int}(B)}}, \hat{Y}'_{\mathcal{E}_{\text{int}(B)}})$$

in our coupling.

These considerations imply that

$$\begin{aligned} \mathbb{Q}(X_W = X'_W) &\geq \mathbb{Q}(T) \\ &\geq 1 - \varepsilon \end{aligned}$$

which concludes the proof as noted above. $\square$

The proof of Proposition 2.9 is an easier application of the concept that the existence of a (quasi-)closed barrier “blocks the information from outside”. Since the proof is virtually the same as the proof of Proposition 3.7 in [18], i.e., the analogous statement for the DaC(1) model, we will just sketch it for the reader’s convenience.

Proof sketch of Proposition 2.9. Fix $W \subset \mathbb{Z}^d$ and $\sigma \in S^{\mathbb{Z}^d \setminus W}$ such that none of the spins in σ percolates. By the assumption (4), this is true for almost every spin configuration. Let $W' \subset \mathbb{Z}^d \setminus W$ be the union of all spin components in $\sigma_{\mathbb{Z}^d \setminus W}$ that intersect the vertex boundary ∂W . Since there is no infinite spin component in σ , we have that W' is a finite set, hence the edge boundary $B = \Delta(W \cup W')$ is a closed barrier. Therefore, it follows from Lemma 3.7 that the conditional distribution of $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $K_{\mathbb{Z}^d \setminus W}^\sigma$ is $\mu_{p,q,(a_1,\dots,a_s)}^{\text{int}(B)}$ conditioned on $\{\xi \in \Omega_C^{\text{int}(B)} : \xi_{W'} = \sigma_{W'}\}$.

Now recall the definition of $\partial_n W$, the n -neighbourhood of W , in Section 2.2. If k is so large that $W' \subset \partial_{k-1} W$, and $\sigma' \in S^{\mathbb{Z}^d \setminus W}$ is such that $\sigma'_{\partial_k W} = \sigma_{\partial_k W}$, then it is clear by the same argument that the conditional distribution of $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$ given $K_{\mathbb{Z}^d \setminus W}^{\sigma'}$ is $\mu_{p,q,(a_1,\dots,a_s)}^{\text{int}(B)}$ conditioned on $\{\xi \in \Omega_C^{\text{int}(B)} : \xi_{W'} = \sigma'_{W'}\}$. Since $\sigma_{W'} = \sigma'_{W'}$, the above conditions are the same, therefore for any $\kappa \in S^W$, we have that

$$|\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^\sigma) - \mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}(K_W^\kappa \mid K_{\mathbb{Z}^d \setminus W}^{\sigma'})| = 0.$$

This proves almost sure quasilocality of $\mu_{p,q,(a_1,\dots,a_s)}^{\mathbb{Z}^d}$. $\square$

Acknowledgements I am grateful to Federico Camia and Ronald Meester for a careful reading of this paper and numerous suggestions, and to Aernout van Enter for drawing my attention to [4], and other valuable comments.

References

- [1] A. Bálint, F. Camia, R. Meester, Sharp phase transition and critical behaviour in 2D divide and colour models, *Stochastic Process. Appl.* **119**, 937–965 (2009).
- [2] A. Bálint, F. Camia, R. Meester, The high-temperature Ising model on the triangular lattice is a critical percolation model, preprint available at arXiv:0806.3020v1 [math.PR] (2008).
- [3] D.J. Barsky, G.R. Grimmett, C.M. Newman, Percolation in half-spaces: equality of critical densities and continuity of the percolation probability, *Probab. Th. Relat. Fields* **90**, 111–148 (1991).
- [4] T. Bodineau, Slab percolation for the Ising model, *Probab. Th. Relat. Fields* **132**, 83–118 (2005).
- [5] A.C.D. van Enter, R. Fernández, A.D. Sokal, Renormalization transformations in the vicinity of first-order phase transitions: What can and cannot go wrong, *Phys. Rev. Lett.* **66**, 3253–3256 (1991).
- [6] A.C.D. van Enter, R. Fernández, A.D. Sokal, Regularity properties of position-space renormalization group transformations: Scope and limitations of Gibbsian theory, *J. Statist. Phys.* **72**, 879–1167 (1993).
- [7] A.C.D. van Enter, C. Maes, R.H. Schonmann, S. Shlosman, The Griffiths singularity random field, *Amer. Math. Soc. Transl. Ser. 2* **198**, 51–58 (2000).
- [8] C.M. Fortuin, P.W. Kasteleyn, J. Ginibre, Correlation inequalities on some partially ordered sets, *Comm. Math. Phys.* **22**, 89–103 (1971).
- [9] O. Garet, Limit theorems for the painting of graphs by clusters, *ESAIM Probab. Statist.* **5**, 105–118 (2001).
- [10] H.-O. Georgii, *Gibbs Measures and Phase Transitions*, de Gruyter Studies in Mathematics **9**, Walter de Gruyter, Berlin (1988).

- [11] H.-O. Georgii, O. Häggström, C. Maes, The random geometry of equilibrium phases, in *Phase Transitions and Critical Phenomena* **18** (C. Domb and J.L. Lebowitz, eds), pp. 1–142, Academic Press, London (2001).
- [12] B. Graham, G. Grimmett, Sharp thresholds for the random-cluster and Ising models, preprint available at arXiv:0903.1501v1 [math.PR] (2009).
- [13] G. Grimmett, The stochastic random-cluster process and the uniqueness of random-cluster measures, *Ann. Probab.* **23**, 1461–1510 (1995).
- [14] G. Grimmett, *Percolation* (2nd ed.), Springer-Verlag, Berlin (1999).
- [15] G. Grimmett, *The random-cluster model*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] **333**, Springer-Verlag, Berlin (2006).
- [16] O. Häggström, Random-cluster representations in the study of phase transitions, *Markov Proc. Relat. Fields* **4**, 275–321 (1998).
- [17] O. Häggström, Positive correlations in the fuzzy Potts model, *Ann. Appl. Probab.* **9**, 1149–1159 (1999).
- [18] O. Häggström, Coloring percolation clusters at random, *Stochastic Process. Appl.* **96**, 213–242 (2001).
- [19] O. Häggström, Is the fuzzy Potts model Gibbsian?, *Ann. Inst. H. Poincaré, Probab. Statist.* **39**, 891–917 (2003).
- [20] C. Külske, (Non-)Gibbsianness and phase transitions in random lattice spin models, *Markov Proc. Relat. Fields* **5**, 357–383 (1999).
- [21] C. Külske, Weakly Gibbsian representations for joint measures of quenched lattice spin models, *Probab. Th. Relat. Fields* **119**, 1–30 (2001).
- [22] J. Kahn, N. Weininger, Positive association in the fractional fuzzy Potts model, *Ann. Probab.* **35**, 2038–2043 (2007).
- [23] C. Maes, K. Vande Velde, The fuzzy Potts model, *J. Phys. A* **28**, 4261–4271 (1995).
- [24] C.M. Newman, L.S. Schulman, Infinite clusters in percolation models, *J. Stat. Phys.* **26**, 613–628 (1981).
- [25] Á. Pisztora, Surface order large deviations for Ising, Potts and percolation models, *Probab. Th. Relat. Fields* **104**, 427–466 (1996).

- [26] V. Strassen, The existence of probability measures with given marginals,
Ann. Math. Statist. **36**, 432–439 (1965).